



WHAT YOU'RE UP AGAINST TREATING GASTRIC ULCERS.



BIOEQUIVALENT

Gastrobim is the **first and only** FDA-approved bioequivalent to GASTROGARD® (omeprazole) under ANADA # 200-842



APPLE FLAVOR

Improve and heal gastric ulcers with an **apple-flavored** formulation



TRUSTED GLOBALLY

The **fastest growing** omeprazole paste globally with more than **one million** doses sold since 2024

Same trusted treatment. Less cost. That's Gastrobim.



PRODUCT	FLAVOR	CONCENTRATION	PASTE	LABELED FOR PREVENTION	BODYWEIGHT TREATED PER SYRINGE	FDA APPROVAL
Gastrobim™ <i>(omeprazole)</i>	Apple	370 mg/g	✓	✓	1250 lbs.	ANADA # 200-842
GASTROGARD™ <i>(omeprazole)</i>	Cinnamon	370 mg/g	✓	✓	1250 lbs.	NADA # 141-123


BIOEQUIVALENCE:

Gastrobim™ is bioequivalent to the pioneer product, GASTROGARD®.

Bioequivalence as defined by FDA is the **absence of a difference** (within pre-defined acceptance criteria) in the **bioavailability** of the active pharmaceutical ingredient (API) or its metabolite(s) at the site of action when administered at the same molar dose **under similar conditions**, in an appropriately designed study.

Following this process, the **conclusion of bioequivalence was confirmed** by the FDA and therefore GASTROGARD® and Gastrobim™ have been accepted as **"interchangeable in a clinical setting"**.

SCAN THE QR CODE TO READ THE FDA DESCRIPTION OF BIOEQUIVALENCE



INDICATIONS:

TREATMENT OF GASTRIC ULCERS:

Gastrobim™ should be administered orally once-a-day for 4 weeks at the recommended dosage of 1.8 mg omeprazole/lb body weight (4 mg/kg).

PREVENTION OF RECURRENCE OF GASTRIC ULCERS:

Continue treatment for at least an additional 4 weeks by administering Gastrobim™ at the recommended daily maintenance dose of 0.9 mg/lb (2 mg/kg).

PRODUCT INFORMATION:

ITEM NO.	PACK SIZE	CASE SIZE
1GAS002	6.15 g	72

For more information and frequently asked questions about Gastrobim, visit www.gastrobim.com/FAQs



IMPORTANT SAFETY INFORMATION: The safety of Gastrobim paste has not been determined in pregnant or lactating mares. For use in horses and foals 4 weeks of age and older. Keep out of reach of children. In case of ingestion, contact a physician. Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.